

# **FOOD SAFETY** is our Passion. Making it **SIMPLE** is our Mission.



# **TECHNICAL DATA BULLETIN**





#### **PRODUCT DESCRIPTION**

**Sani Professional® Hands Instant Sanitizing Wipes** are nonwoven wipes saturated with an ethyl alcohol solution for the antimicrobial cleansing of hands. Solution and towel are fragrance-free and dye-free.

Sani Professional Sani Professional Item # Item # Item # P92084 D33333 P43572

# CHEMICAL COMPOSITION/PRODUCT DATA

Active Ingredients: Alcohol (ethyl alcohol) 70% (by volume)

Inactive Ingredients: Water, Propylene glycol, Glycerin, Aloe barbadensis leaf juice, Tocopheryl acetate (Vitamin E)

30%

TOTAL 100.00%

(Does not include weight of the wipe)



#### **EFFICACY STUDIES**

#### **BACTERIA**

#### **IN-VITRO TIME KILL STUDIES**

**Purpose** – To determine how rapidly and effectively **Sani Professional Hands Instant Sanitizing Wipes** killed a variety of Gram negative and Gram positive microorganisms after a 15-second exposure.

**Methodology** – Fluid from the wipe was expressed aseptically and transferred to sterile incubator tubes. The tubes were subsequently inoculated with the broth culture of each test microorganism containing up to 10<sup>8</sup> CFU/ml. After 15 seconds, the entire inoculated volume of **Sani Professional Hands Instant Sanitizing Wipes** was transferred to neutralizers. Serial dilutions were plated using standard plating techniques and percent reductions for each organism were calculated after incubation.

**Conclusion – Sani Professional Hands Instant Sanitizing Wipes** proved to be effective at killing all 32 microorganisms listed within a 15 second exposure.

Independent Laboratory: Mycoscience Labs, Willington, CT: June 28, 2004

#### **CHART 1: PERCENT REDUCTION AFTER 15-SECOND EXPOSURE**

Microorganism	Classification	ATCC#	% Reduction
Acinetobacter baumanii (multi-drug resistant)	Gram negative rod	19606	>99.999
Aspergillus flavus	fungi (mold)	9643	>99.999
Bacillus megaterium	Gram positive rod	14581	>99.999
Campylobacter jejuni	Gram negative rod	29428	>99.999
Candida albicans	fungi (yeast)	14053	>99.999
Clostridium difficile (vegetative)	Gram positive rod	9689	>99.998
Community Acquired Methicillin Resistant Staphylococcus aureus (CA-MRSA) [NARSA NRS384] [Genotype USA 300]			>99.999
Community Acquired Methicillin Resistant Staphylococcus aureus (CA-MRSA) [NARSA NRS123] [Genotype USA 400]			>99.999
Corynebacterium diptheriae	Gram positive rod	11913	>99.999
Enterobacter aerogenes	Gram negative rod	13048	>99.999
Enterococcus faecium (multi-drug resistant including Vancomycin)	Gram positive cocci	51559	>99.999
Enterococcus faecalis (Vancomycin, Streptomycin, and Gentamicin resistant)	Gram positive cocci	51575	>99.999
Escherichia coli (ESBL producing, multi-drug resistant, derived from clinical isolate, Klebsiella pneumoniae ATCC #14714)	Gram negative rod	BAA-196	>99.999
Escherichia coli (0157:H7)	Gram negative rod	11229	>99.999
Escherichia coli (0111:H8)	Gram negative rod	BAA-184	>99.999
Klebsiella pneumoniae	Gram negative rod	13883	>99.999
Klebsiella pneumoniae (NDM-1 Positive) [CDC 1000527]			>99.999
Klebsiella pneumoniae (carbapenem resistant)	Gram negative rod	BAA-1705	>99.999
Listeria monocytogenes	Gram positive rod	15313	>99.999
Proteus mirabilis	Gram negative rod	7002	>99.999
Proteus hauseri (vulgaris)	Gram negative rod	13315	>99.999
Pseudomonas aeruginosa	Gram negative rod	15442	>99.999
Salmonella choleraesuis serotype typhimurium	Gram negative rod	14028	>99.999
Serratia marcescens	Gram negative rod	14756	>99.999
Shigella sonnei	Gram negative rod	11060	>99.999
Staphylococcus aureus (MRSA)	Gram positive rod	33591	>99.999
Staphylococcus aureus (MRSA, Vancomycin tolerant)	Gram positive rod	700788	>99.999
Staphylococcus epidermidis	Gram positive cocci	12228	>99.999
Streptococcus pneumoniae	Gram positive cocci	33400	>99.999
Streptococcus pyogenes	Gram positive cocci	19615	>99.999
Trichophyton interdigitale (formerly mentagrophytes)	fungi (mold)	9533	>99.999
Vibrio parahaemolyticus	Gram negative rod	17802	>99.999





# **KILLS TOP FOODBORNE PATHOGENS**

**Sani Professional Hands Instant Sanitizing Wipes** have been proven effective against the following foodborne pathogens:

SOURCES:	
Raw or undercooked poultry, unpasteurized (raw) milk, contaminated drinking water	
Undercooked ground beef, unpasteurized (raw) milk or juice, soft cheeses made from raw milk, and raw fruits and vegetables (such as sprouts)	
Raw produce, contaminated drinking water, uncooked foods and cooked foods that are not reheated after contact with an infected food handler	
Ready-to-eat deli meats and hot dogs; refrigerated pâtés or meat spreads; unpasteurized (raw) milk and dairy products; soft, unpasteurized cheeses (e.g., queso fresco, Feta, Brie, Camembert); refrigerated smoked seafood; raw sprouts	
Eggs, poultry, meat, unpasteurized (raw) milk or juice, cheese, contaminated raw fruits and vegetables	
Undercooked or raw seafood, such as shellfish (especially oysters)	

EFFECTIVE AGAINST

99.99%
OF MOST COMMON BACTERIA ON HANDS



\*99.999% reduction of the microorganisms achieved within 15 seconds Source: www.cdc.gov/foodsafety



#### ADDITIONAL INFORMATION AND TESTING

#### **SAFETY**

# **Repeated Insult Patch Test**

Purpose – To determine the dermal irritation and sensitization potential of Sani Professional Hands Instant Sanitizing Wipes.

**Methodology** – Study was conducted using 216 subjects. The induction phase involved repeated exposure of the product at the same site on each subject three times a week for a total of nine applications. Ten to 14 days after induction, a challenge patch was applied to a virgin site on each subject for 24 hours. After 24 hours, the patch was removed and the site was evaluated for dermal irritation.

**Conclusion – Sani Professional Hands Instant Sanitizing Wipes** demonstrated minimal or no reaction which would cause dermal irritation or sensitization.

Independent Laboratory: Clinical Research Laboratories, Piscataway, NJ: June 11, 2004

#### **SAFETY IN USE**

# **Modified Safety In Use Test**

**Purpose –** To evaluate the dermal irritation potential of **Sani Professional Hands Instant Sanitizing Wipes** under exaggerated use conditions following 25 repeated uses.

**Methodology** – A total of 25 human subjects completed the study. Each subject used one wipe on both hands for approximately 30 seconds. This was repeated 25 times with 5-minute intervals between uses. Subjects hands were evaluated at the end of 25 uses.

**Conclusion – Sani Professional Hands Instant Sanitizing Wipes** did not demonstrate any potential for eliciting dermal irritation in any of the 25 human subjects.

Independent Laboratory: Clinical Research Laboratories, Piscataway, NJ: May 13, 2004

#### FOODSERVICE INFORMATION AND TESTING

# **FDA Food Code Compliant**

Meets the Food and Drug Administration (FDA) Food Code, Section 2-301.16.

# NSF Nonfood Compounds Registration #151433 Category E3

This product is acceptable for use as a hand sanitizing product in and around food processing areas.

# Kills Top Foodborne Pathogens

Tested 99.999% effective in 15 seconds against *Campylobacter jejuni, Escherichia coli* (0157:H7), *Listeria monocytogenes, Salmonella choleraesuis serotype typhimurium, Shigella sonnei* and *Vibrio parahaemolyticus*.

# OTHER INFORMATION AND TESTING

#### **Skin Moisturization**

Sani Professional Hands Instant Sanitizing Wipes contains soothing Aloe and Vitamin E.

# OSHA Bloodborne Pathogen Standard 29 CFR Part 1910.1030

Meets the specific handwashing standard 1910.1030 (d)(2)(iv).

#### **Glove Use**

It is recommended to allow hands to dry completely after using **Sani Professional Hands Instant Sanitizing Wipes** prior to applying gloves.

#### **Shelf Life**

Stability testing was conducted for purposes of establishing an expiration date for the unopened product. Current stability data supports a two-year expiration period from the date of bulk liquid manufacture.

Data on file

# **PRECAUTIONARY STATEMENTS**

Flammable, keep away from fire or flame.

For external use only.

Do not use in or contact the eyes.

Discontinue use if irritation and redness develop.

If condition persists for more than 72 hours consult a physician.



61041v8

Made in USA with domestic and imported material